

**Quão enviesadas são medidas de mobilidade
intergeracional de educação?
Orientação prática de um modelo de simulação**

Marcelo Cardona & Sam Jones

27 Outubro 2021

Overview

- Focus is on the measurement of intergenerational (educational) mobility
- Bewildering range of empirical choices face applied researchers
- Evidence on bias associated with different choices is scattered and incomplete
- Address this gap using a flexible simulation framework
- Provide some practical guidance
- Application to Mozambique

Agenda


- 1** Measuring mobility
- 2** Simulation framework
- 3** Results: stylized low education case
- 4** Results: generalized case
- 5** Mozambique application
- 6** Conclusion

(1) Measuring mobility

Broad terrain

- Broad sense, IGM refers to the degree to which advantages and disadvantages of individuals persist across generations (e.g., Great Gatsby curve) – dimension of social justice
- Relevant in multiple domains, but education often a primary focus in developing countries (as robust predictor of well-being and measured directly)
- No single accepted definition: “IGM is a complex concept and may mean different things to different people ... A natural consequence ... is that there is no consensus on how IGM should be measured, so many indices are available to an applied researcher.” (Savegnago, 2016: 386)

Stylized linear example



Challenge 1: Multiple metrics

Define: c = child outcome; p = parental outcome

| Class | Focus | Example |
|--------------|------------|---------------------------------------|
| Heritability | Slope | $\text{Correl}(c,p)$ |
| Performance | Distance | $E(c p) - p$ |
| | Transition | $\Pr(c > p p \leq p^*)$ |
| Distribution | Inequality | $\text{Gini}(c + p) / \text{Gini}(p)$ |

Distinctions between:

- Parametric (regression) vs. non-parametric approaches (e.g., transition matrices)
- Number of free parameters (to be estimated)

Challenge 2: Multiple transforms

Monotonic data transformations often used to remove (some) nuisance parameters and facilitate comparison/interpretation

| Transform | Mean | Variance |
|-----------------------------|-------|----------|
| None | μ | σ |
| Variance stabilization | μ | 1 |
| Mean-variance stabilization | 0 | 1 |
| Rank | 1/2 | 1/12 |

Issues:

- Applied separately or jointly?
- Sensitivity to reference distribution(s) & outliers

Challenge 3: Multiple data issues

Few parent-child matched administrative datasets exist (outside Scandinavia).

Co-resident, self-reported data is commonplace (e.g., census).

Three generic issues → alternative corrections:

| Problem | Poss. corrections | Example |
|----------------------|----------------------|---|
| Incomplete education | Sample restrictions | Child ≥ 18 |
| Reporting error | Ranking | Percentile |
| Missing co-residents | Prediction, bounding | $c - \epsilon \leq p \leq c + \epsilon$ |

⇒ expect to generate bias!

(2) Simulation framework

The problem

Open question: **exactly *how* biased are the different measures of IGM** (metrics \times transforms)?

- are all measures equally biased?
- do conventional empirical corrections matter?
- is bias more severe in different contexts?

Difficult to answer analytically:

- Different sources of bias *may* offset one another [see paper]
- Various measures are non-linear (in expectation)

So, let's run some scenarios!

Basic set-up

Extension of generalized errors-in-variables approach (c.f., Nybom and Stuhler, 2017), to account for systematic missing observations and incomplete education:

$$c_{ij}^* = \alpha_j \bar{p}^* + \beta_j p_i^* + \theta_0 \varepsilon_{0ij} \quad (1a)$$

$$c_{ij} = c_0 + (1 + \lambda_c) c_{ij}^* - \delta m_i^* + \nu_i \quad (1b)$$

$$p_i = p_0 + (1 + \lambda_p) p_i^* + \mu_i \quad (1c)$$

$$\nu_i = \theta_1 \varepsilon_{1i}, \quad \mu_i = \gamma \nu_i + \theta_2 \varepsilon_{2i} \quad (1d)$$

$$\varepsilon_{1ij} \sim \mathcal{N}_I^u(0, \sqrt{1 - \beta_j^2} \cdot \sigma_{p^*}), \quad \varepsilon_{1i}, \varepsilon_{2i} \sim \mathcal{N}_I^u(0, 1) \quad (1e)$$

where i are individuals and j indexes seen vs unseen groups;

\mathcal{N}_I^u is a truncated normal distribution; and

$\theta_0, \theta_1, \theta_2 \geq 0$ are scaling parameters.

Metrics × transforms × corrections

| Metric | Expression | Description |
|--------------------|---|----------------------------|
| Non-Herit. | $1 - \beta$ | Reverse of slope |
| Out-perform. | $\hat{\alpha} + p_{25}(\hat{\beta} - 1)$ | Exp. diff at 25th pc |
| Inequality | $1 - g(c + p)/g(c)$ | Fall in inequality |
| Move sh. | $N^{-1} \sum_N \mathbf{1}(c - p > \pi)$ | Share moved by $> \pi$ |
| Move sh. (+) | $N^{-1} \sum_N \mathbf{1}(c - p > \pi)$ | Share moved upward |
| Move sh. (++) | $N^{-1} \sum_N \mathbf{1}(c - p > \pi) \mid p < p_{50}$ | Conditional share moved up |
| Wgt. move sh. | $N^{-1} \sum_N c - p / \max(c - p)$ | Weighted share moved |
| Wgt. move sh. (+) | as above $\cdot \mathbf{1}(c^r > p^r)$ | Weighted share moved up |
| Wgt. move sh. (++) | as above $\mid p < p_{50}$ | Conditional share moved up |

Metrics × transforms × corrections

| Transform | Description |
|-----------|--------------------------------------|
| None | - |
| Var. | Variance stabilization (separate) |
| Ref. | Percentiles of parental distribution |
| Rank | Percentiles (separate) |

| Correction | Description |
|------------|--|
| None | - |
| Enroll | Only children NOT enrolled in school |
| Age | Only children with age such that $c > p$ |
| Predict | Predict missing p from c (age-corrected) |

Implementation steps

- 1 Calibrate parameters of true DGP (e.g., ρ^* , α_j , β_j)
- 2 Set assumptions for measurement error structure (e.g., θ_1 , θ_2 , ρ_0 , c_0 , λ_c , λ_p , γ , δ)
- 3 Draw stochastic variables (error terms, iid)
- 4 Simulate full dataset
- 5 Impose corrections (to observed data) and transforms
- 6 Calculate mobility metrics (true vs observed)

Note: step 6 can be bootstrapped to verify error distributions.

Measures of bias

Here for β , but can be any chosen final metric:

$$\text{MB}(\beta) = \frac{1}{N} \sum_{n=1}^N (\hat{\beta}_n - \beta_n) \quad (2a)$$

$$\text{MFB}(\beta) = \frac{1}{N} \sum_{n=1}^N \frac{2(\hat{\beta}_n - \beta_n)}{(|\hat{\beta}_n| + |\beta_n|)} \quad (2b)$$

$$\text{MAFB}(\beta) = \frac{1}{N} \sum_{n=1}^N \frac{2(|\hat{\beta}_n - \beta_n|)}{(|\hat{\beta}_n| + |\beta_n|)} \quad (2c)$$

where $n = \{1, \dots, N\}$ indexes simulation iterations (for a given scenario).

(3) Results: stylized low education case


Stylized low education case

Calibrate model to Mozambique and consider 6 distinct measurement error scenarios.


Table: Mean outcomes (all 6 scenarios \times 50 iterations = 300 obs.)

| Metric \downarrow // Transform \rightarrow | True | Estimated bias (no correction) | | | |
|--|------|--------------------------------|-------|-------|-------|
| | None | None | Var | Ref. | Rank |
| Non-Herit. | 0.60 | 0.05 | 0.06 | 0.03 | 0.04 |
| Out-Perform. | 0.17 | 0.01 | 0.01 | -0.01 | 0.01 |
| Inequality | 0.54 | -0.00 | -0.00 | -0.01 | 0.02 |
| Move sh. | 0.89 | -0.06 | 0.00 | 0.01 | 0.00 |
| Move sh. (+) | 0.61 | -0.05 | -0.02 | -0.02 | -0.00 |
| Move sh. (++) | 0.79 | -0.05 | -0.01 | -0.03 | -0.01 |
| Wgt. m. sh. | 0.22 | 0.00 | 0.01 | 0.00 | 0.01 |
| Wgt. m. sh. (+) | 0.16 | -0.01 | 0.00 | -0.01 | 0.00 |
| Wgt. m. sh. (++) | 0.22 | -0.00 | 0.01 | -0.02 | -0.00 |


All scenarios, no corrections (metrics 1-3)




All scenarios, no corrections (metrics 4-6)




All scenarios, no corrections (metrics 7-9)




All scenarios, ref. transform (metrics 1-3)



All scenarios, ref. transform (metrics 4-6)



All scenarios, ref. transform (metrics 7-9)




(4) Results: generalized case

Generalized scenarios, no corrections


Run 300 separate draws of *both* DGP and measurement error structure (jointly).

On each draw look at bias associated with all combinations of transforms and corrections.

Non-heritability



Move sh. (++)



| | Non-herit | | Out-Perf. | | Move % (++) | |
|--------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|
| Constant | 0.150*** (0.004) | 0.150*** (0.004) | -0.065*** (0.006) | -0.065*** (0.006) | -0.030*** (0.003) | -0.030*** (0.003) |
| Var. trans. | 0.033*** (0.005) | 0.033*** (0.005) | 0.088*** (0.006) | 0.088*** (0.006) | 0.022*** (0.003) | 0.022*** (0.003) |
| × α | | -0.035*** (0.006) | | -0.037*** (0.008) | | 0.006* (0.003) |
| × β | | 0.003 (0.006) | | -0.013** (0.006) | | -0.002 (0.003) |
| × \bar{p} | | -0.012** (0.005) | | 0.057*** (0.008) | | 0.022*** (0.003) |
| Ref. trans. | -0.051*** (0.004) | -0.051*** (0.003) | 0.011 (0.006) | 0.011* (0.006) | 0.002 (0.003) | 0.002 (0.003) |
| × α | | -0.045*** (0.004) | | 0.023** (0.009) | | 0.014*** (0.003) |
| × β | | -0.046*** (0.004) | | -0.001 (0.006) | | 0.001 (0.003) |
| × \bar{p} | | 0.027*** (0.004) | | 0.015* (0.009) | | -0.002 (0.003) |
| Rank. trans. | 0.055*** (0.004) | 0.055*** (0.004) | 0.232*** (0.008) | 0.232*** (0.008) | 0.081*** (0.004) | 0.081*** (0.004) |
| × α | | -0.001 (0.005) | | 0.032*** (0.011) | | 0.019*** (0.004) |
| × β | | 0.025*** (0.005) | | 0.046*** (0.008) | | 0.018*** (0.004) |
| × \bar{p} | | 0.004 (0.004) | | 0.007 (0.010) | | -0.015*** (0.004) |
| Enroll corr. | 0.082*** (0.006) | 0.082*** (0.006) | -0.114*** (0.010) | -0.115*** (0.010) | -0.043*** (0.005) | -0.043*** (0.005) |
| Age corr. | -0.037*** (0.005) | -0.037*** (0.005) | 0.023*** (0.007) | 0.023*** (0.007) | 0.004 (0.003) | 0.004 (0.003) |
| Fit corr. | -0.020*** (0.006) | -0.020*** (0.006) | -0.076*** (0.009) | -0.076*** (0.009) | -0.027*** (0.004) | -0.027*** (0.004) |
| Obs. | 4,800 | 4,800 | 4,800 | 4,800 | 4,800 | 4,800 |
| R2 (adj.) | 0.644 | 0.681 | 0.598 | 0.615 | 0.516 | 0.537 |
| RMSE | 0.107 | 0.102 | 0.172 | 0.168 | 0.075 | 0.073 |

| | Non-herit | | Out-Perf. | | Move % (++) | |
|--------------|----------------------|----------------------|----------------------|----------------------|----------------------|----------------------|
| Constant | 0.160*** (0.003) | 0.160*** (0.003) | 0.227*** (0.005) | 0.227*** (0.005) | 0.093*** (0.002) | 0.093*** (0.002) |
| Var. trans. | 0.047*** (0.004) | 0.047*** (0.004) | -0.104*** (0.005) | -0.104*** (0.005) | -0.043*** (0.002) | -0.043*** (0.002) |
| × α | | -0.010** (0.005) | | 0.045*** (0.006) | | 0.010*** (0.003) |
| × β | | 0.016*** (0.005) | | 0.023*** (0.005) | | 0.003 (0.002) |
| × \bar{p} | | -0.006 (0.004) | | -0.037*** (0.006) | | -0.022*** (0.003) |
| Ref. trans. | -0.061*** (0.003) | -0.061*** (0.003) | -0.048*** (0.005) | -0.048*** (0.005) | -0.031*** (0.002) | -0.031*** (0.002) |
| × α | | -0.043*** (0.004) | | 0.003 (0.007) | | 0.005** (0.003) |
| × β | | -0.051*** (0.004) | | -0.002 (0.005) | | -0.006*** (0.002) |
| × \bar{p} | | 0.025*** (0.004) | | 0.007 (0.007) | | -0.003 (0.003) |
| Rank. trans. | 0.042*** (0.004) | 0.042*** (0.003) | -0.032*** (0.007) | -0.032*** (0.006) | -0.015*** (0.003) | -0.015*** (0.003) |
| × α | | -0.000 (0.004) | | 0.117*** (0.008) | | 0.035*** (0.003) |
| × β | | 0.018*** (0.004) | | 0.107*** (0.006) | | 0.018*** (0.003) |
| × \bar{p} | | 0.003 (0.004) | | -0.056*** (0.007) | | -0.027*** (0.003) |
| Enroll corr. | 0.075*** (0.005) | 0.075*** (0.005) | 0.028*** (0.009) | 0.028*** (0.008) | 0.019*** (0.003) | 0.019*** (0.003) |
| Age corr. | -0.014*** (0.004) | -0.014*** (0.004) | -0.009 (0.006) | -0.009* (0.005) | -0.003 (0.002) | -0.003 (0.002) |
| Fit corr. | -0.014*** (0.005) | -0.014*** (0.004) | 0.048*** (0.008) | 0.048*** (0.007) | 0.015*** (0.003) | 0.015*** (0.003) |
| Obs. | 4,800 | 4,800 | 4,800 | 4,800 | 4,800 | 4,800 |
| R2 (adj.) | 0.713 | 0.752 | 0.393 | 0.504 | 0.428 | 0.475 |
| RMSE | 0.089 | 0.083 | 0.145 | 0.131 | 0.058 | 0.055 |

Main lessons

- 1 Non-heritability coefficient ($1 - \beta$) upward biased
- 2 Lower bias of non-parametric measures (e.g., % up)
- 3 Data transforms *often* reduce bias (mean & absolute)
- 4 Additional corrections do not generally reduce bias
- 5 BUT no single "always-lowest-bias" measure – characteristics of the case matter (shapes of true CDFs)

Recommendation


Use a *calibrated* simulation model to estimate upper/lower bounds on bias for a plausible range of 'true' models.
e.g., 95% of time bias is within $|\varepsilon|$.

(5) Mozambique application

Median years of education by province

| Province | Child's education | | | Parent's education | | |
|-----------------|-------------------|------|------|--------------------|------|------|
| | 1997 | 2007 | 2017 | 1997 | 2007 | 2017 |
| Niassa | 2 | 3 | 5 | 2 | 3 | 6 |
| Cabo Delgado | 2 | 3 | 4 | 2 | 3 | 5 |
| Nampula | 2 | 3 | 5 | 3 | 4 | 5 |
| Zambezia | 2 | 3 | 4 | 2 | 3 | 5 |
| Tete | 2 | 3 | 5 | 2 | 3 | 5 |
| Manica | 2 | 4 | 5 | 2 | 4 | 6 |
| Sofala | 3 | 4 | 5 | 3 | 4 | 6 |
| Inhambane | 3 | 4 | 4 | 3 | 4 | 5 |
| Gaza | 3 | 4 | 4 | 3 | 4 | 4 |
| Maputo-province | 4 | 6 | 6 | 4 | 6 | 7 |
| Maputo-city | 5 | 7 | 7 | 6 | 7 | 7 |


Mean education of children vs. parents, 1997-2017



Measures of IGM using a reference distribution

| Province | Upwards-mobility | | | Out-performance | | |
|-----------------|------------------|------------------|------------------|------------------|------------------|------------------|
| | 1997 | 2007 | 2017 | 1997 | 2007 | 2017 |
| Niassa | 0.372 (0.002) | 0.601 (0.002) | 0.395 (0.002) | 0.223 (0.001) | 0.264 (0.001) | 0.138 (0.001) |
| Cabo Delgado | 0.363 (0.002) | 0.632 (0.002) | 0.336 (0.002) | 0.230 (0.001) | 0.281 (0.001) | 0.118 (0.001) |
| Nampula | 0.392 (0.001) | 0.637 (0.001) | 0.352 (0.001) | 0.201 (0.001) | 0.248 (0.001) | 0.126 (0.001) |
| Zambezia | 0.420 (0.001) | 0.682 (0.001) | 0.351 (0.001) | 0.251 (0.001) | 0.292 (0.001) | 0.156 (0.001) |
| Tete | 0.387 (0.001) | 0.625 (0.001) | 0.425 (0.001) | 0.250 (0.001) | 0.300 (0.001) | 0.170 (0.001) |
| Manica | 0.462 (0.002) | 0.810 (0.001) | 0.527 (0.002) | 0.256 (0.001) | 0.366 (0.001) | 0.214 (0.001) |
| Sofala | 0.429 (0.002) | 0.766 (0.001) | 0.523 (0.001) | 0.211 (0.001) | 0.317 (0.001) | 0.208 (0.001) |
| Inhambane | 0.664 (0.002) | 0.860 (0.001) | 0.736 (0.002) | 0.364 (0.001) | 0.427 (0.001) | 0.329 (0.001) |
| Gaza | 0.656 (0.002) | 0.854 (0.001) | 0.685 (0.002) | 0.352 (0.001) | 0.409 (0.001) | 0.319 (0.001) |
| Maputo-Province | 0.729 (0.002) | 0.750 (0.001) | 0.694 (0.001) | 0.298 (0.001) | 0.351 (0.001) | 0.254 (0.001) |
| Maputo-City | 0.548 (0.001) | 0.806 (0.001) | 0.780 (0.002) | 0.274 (0.001) | 0.338 (0.001) | 0.271 (0.001) |
| Mozambique | 0.483 (0.000) | 0.719 (0.000) | 0.493 (0.000) | 0.243 (0.000) | 0.300 (0.000) | 0.200 (0.000) |

Upward mobility (++) , using a ref. distribution



(6) Conclusion

Conclusion

- Multiple measures of IGM, various sources of bias
- Set-up a (calibrated) simulation framework
- Bias largest for heritability coefficient BUT no all round "best" measure
- In Mozambique, simulations suggest out-performance or upward mobility generally least biased (+/- 5%)
- Census data indicates large spatial heterogeneity in IGM in Mozambique
- Weakening of IGM since 2007, esp. in North